

*The 12th International Symposium on Biogeochemistry of Wetlands in  
Coral Springs, Florida, USA, April 23-26, 2018*

# **Methane Emissions from the Stems of Living Trees in Upland Forests**

**Zhi-Ping Wang**

**Institute of Botany, Chinese Academy of  
Sciences, Beijing**






# Acknowledgements

Pat Megonigal  
Sunitha Pangala

科技  
部

Ministry of Science and Technology



中国科学院植物研究所  
INSTITUTE OF BOTANY, THE CHINESE ACADEMY OF SCIENCES






# Outline

---


- 1. Do plant-based CH<sub>4</sub> emissions constitute a distinct source?**
- 2. CH<sub>4</sub> production / emissions in / from the heartwood / stems of living trees**
- 3. Factors controlling CH<sub>4</sub> production in heartwood**
- 4. Perspective: Novel and large source?**

# Are plant-based $\text{CH}_4$ emissions a distinct source?



*Carmichael et al. 2014*

Plant-based emission of 32–143 Tg CH<sub>4</sub> yr<sup>-1</sup> (*Carmichael et al., 2014*)?







# Forest Wetlands versus Upland Forests

5%

Area

95%








# Upland Forests



# Outline

## 2. CH<sub>4</sub> production / emissions in / from the heartwood / stems of living trees



Research

## Methane emissions from the trunks of living trees on upland soils


Zhi-Ping Wang<sup>1,2</sup>, Qian Gu<sup>1</sup>, Feng-Dan Deng<sup>1,3</sup>, Jian-Hui Huang<sup>1</sup>, J. Patrick Megonigal<sup>4</sup>, Qiang Yu<sup>2</sup>,  
Xiao-Tao Lü<sup>2</sup>, Ling-Hao Li<sup>1</sup>, Scott Chang<sup>5</sup>, Yun-Hai Zhang<sup>1</sup>, Jin-Chao Feng<sup>6</sup> and Xing-Guo Han<sup>1,2</sup>

<sup>1</sup>State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Nanxincun 20, Xiangshan, Beijing 100093, China; <sup>2</sup>State Key Laboratory of Forest and Soil Ecology, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110164, China; <sup>3</sup>University of Chinese Academy of Sciences, Yuquanlu, Beijing 100049, China;

<sup>4</sup>Smithsonian Environmental Research Center, PO Box 28, Edgewater, MD 21037-0028, USA; <sup>5</sup>Department of Renewable Resources, University of Alberta, Edmonton, T6G 2E3 Alberta,

Canada; <sup>6</sup>Institute of Desertification Studies, Chinese Academy of Forestry, Beijing 100091, China









The experimental layout in the Beijing Forest Ecosystem Station (a), the upper plot (b), the lower plot (c), and chambers (d).

July 2014 – July 2015














CH<sub>4</sub> production in the heartwoods of living *P. davidiana* in a small terrace in the Xiaolongmen Forest Farm.

Position	Average age (y)	Trunk diameter (cm)	Heartwood		
			Water content (%)	CH <sub>4</sub> concentration (μL L <sup>-1</sup> )	CH <sub>4</sub> production (ng gdw <sup>-1</sup> h <sup>-1</sup> )
118°44'6.8"E, 31°57'3.5"N 1150 m above sea level	20	47.3 (5.7)	68.5 (3.0)	23.6 (20.0) × 10 <sup>4</sup>	81.64 (63.09)

Value is Mean (SD), *n* = 5 for trunks.

Wood materials were sampled in August 9, 2015.






## Previous studies



## Our study







# CH<sub>4</sub> flux calculations and CH<sub>4</sub> budget estimates

**Table 1** Annual budget of CH<sub>4</sub> in the forest ecosystem

Component	Jul 2014	Aug	Sep	Oct	Nov	Dec	Jan 2015	Feb	Mar	Apr	May	Jun	Jul	Annual	
CH <sub>4</sub> flux (μg trunk <sup>-1</sup> h <sup>-1</sup> for tree or μg m <sup>-2</sup> h <sup>-1</sup> for soil)															
Tree	<i>Populus davidiana</i>														
	Trunk I	1345.3	1318.1	517.9	530.8	309.9	n.a.		312.3	606.2	922.4	1240.1	1083.4		
	Trunk II	1125.0	1118.8	342.2	398.9	215.6			309.7	502.4	819.8	1137.4	841.4		
	Trunk III	1185.7	1170.7	411.1	437.7	276.2			310.2	525.9	837.4	1160.6	936.0		
	Twig and leaf		n.a.	n.a.											
	<i>Carya cathayensis</i>														
	<i>Larix gmelinii</i>														
Soil	-74.5	-73.2	-60.3	-50.6	-47.0	-12.1			-19.3	-39.2	-58.2	-79.5	-57.2		
Plot-wide CH <sub>4</sub> (g plot <sup>-1</sup> )															
Tree	Mean	76.2	75.2	25.6	28.5	16.2			19.4	33.0	53.7	71.3	59.6	390.7	
	Range	70.3~84.1	69.9~82.4	20.7~31.3	24.9~33.2	13.0~18.7			19.4~19.5	30.4~36.7	51.2~57.6	68.8~75.0	52.6~67.1	359.8~430.3	
Soil		-88.7	-87.2	-69.5	-60.2	-54.1	-14.4	-14.4	-13.9	-23.0	-45.1	-69.3	-91.6	-68.1	-621.1

Plot-wide CH<sub>4</sub> were estimated using the parameters of living tree species, such as the 84 trunks of living *P. davidiana* in the lower plot of 1600 m<sup>2</sup> (Tree bases were assumed as zero and not excluded in plot area) and the mean 15.3 m trunk height of *P. davidiana* (see Table S1).

Trunk I, II, and III indicate trunk CH<sub>4</sub> emissions calculated by arithmetic average, logarithm function, and power function, respectively.


Annual CH<sub>4</sub> is the sum of those in months; CH<sub>4</sub> in July is an average of two values in July 2014 and July 2015.

The CH<sub>4</sub> fluxes measured were undetectable and defined as n.a. for not applicable (no data available).

# Traditional hypothesis



# Our result





# Outline

## 3. Factors controlling CH<sub>4</sub> production in heartwood

Tree species with capacity of substantial CH<sub>4</sub> production in heartwood

Temperature

Water content

(*Wang et al., 2017, Journal of Geophysical Research: Biogeosciences*)


Most of tree species with no capacity of substantial CH<sub>4</sub> production in heartwood (*Wang et al., 2017*)

Even if high water content, no CH<sub>4</sub> production!

**Why?**

Secondary metabolites, e.g. carbohydrates, phenolic compounds?  
(Unpublished)

# Field investigation



Mid-temperate

20×20 m<sup>2</sup> plots



Sub-tropical





Warm temperate



Increment borer







The ranges:

R1 < 21.6

21.6 ≤ R2 < 100



100 ≤ R3 < 1,000




1,000 ≤ R4 < 10,000

R5 ≥ 10,000 μL L<sup>-1</sup>

**CH<sub>4</sub> emission?**







Heartwood water content (w/w, %)



Extractive solution of heartwood of a tree species that has no capacity of substantial CH<sub>4</sub> production in heartwood




Adding



Answering why no capacity of substantial CH<sub>4</sub> production


Fresh heartwood of another tree species that has the capacity of substantial CH<sub>4</sub> production in heartwood



## Water-soluble extractives enhanced microbial CH<sub>4</sub> production.


**(a)** Treatments are the 1 g heartwood of *Ulmus davidiana* + 1 mL Deionized Water, + Extractive (1 mL heartwood extractive solution of *Salix matsudana* (no CH<sub>4</sub> production)) and + autoclaved Extractive. **(b)** Treatments are the 1 g heartwood of *U. davidiana* or *P. canadensis* + 1 mL DW and + 1 mL heartwood extractive solution of *P. tabuliformis* (no CH<sub>4</sub> production).





**Ethanol-soluble extractives enhanced microbial CH<sub>4</sub> production, when ethanol's effect was removed.**

Treatments are the 0.8 g fresh heartwood of *Populus davidiana* + 1 mL DW and + 1 mL 100% E extractive solution of *Pinus tabuliformis* (no CH<sub>4</sub> production) evaporated and then 1 mL DW added.



## Acetone-soluble extractives inhibited microbial CH<sub>4</sub> production, when acetone's effect was removed.

Treatments were the 1 g fresh heartwood of *P. canadensis* or *U. davidiana* + 1 mL DW or 1 mL heartwood extractive solution of *P. tabuliformis* (no CH<sub>4</sub> production), the 1 g fresh heartwood of *P. davidiana* + 1 mL DW or + 1 mL heartwood extractive solution of *S. matsudana* (no CH<sub>4</sub> production).



**No substantial CH<sub>4</sub> in the heartwood of most tree species in upland forests, when other variables are appropriate**



**Net effect of secondary metabolites on microbial CH<sub>4</sub> production in heartwood?**

**Inhibition!**

# Summary

- For tree species with capacity of substantial CH<sub>4</sub> production, when temperature was not a limiting factor for CH<sub>4</sub> production in summer and autumn, and thus, most of the CH<sub>4</sub> production may be explained by **water content** in the heartwood of living trees.
- For tree species with no capacity of substantial CH<sub>4</sub> production, net effect of **secondary metabolites** may inhibit microbial CH<sub>4</sub> production in the heartwood.



# Perspective: Novel and large source?

---

- 1, Ecogeography: temporal and spatial distribution, the magnitude of CH<sub>4</sub> emission?*
- 2, CH<sub>4</sub> production mechanism and dynamics?*
- 3, More relationships between CH<sub>4</sub> production/emission and factors?*

**Thank you for attention**

