

Yiyang Kang¹, David A. Kaplan², Michael J. Osland³

¹School of Natural Resources and Environment, University of Florida, Gainesville, Florida, USA

²Engineering School of Sustainable Infrastructure and Environment, University of Florida, Gainesville, Florida, USA



³Wetland and Aquatic Research Center, U.S. Geological Survey, Lafayette, Louisiana, USA

INTRODUCTION

- Alleviated freeze stress induced by the warming climate has resulted in range expansion of mangroves¹.
- Understanding how mangroves respond to freezing by temperature, species, and stratum is critical for predicting future coastal wetlands transformation^{2, 3}.

METHODS


- Gradient study with 12 field sites across northern Florida's Gulf of Mexico coast
- Plot study: Freeze response by temperature, species, and stratum
- Tagged trees: Low temperature thresholds by species



Mangroves in (a) Cedar Key and (b, c) Goose Island, FL before and after the freeze event in 2022. Photo credit: Yiyang Kang

RESULTS


Temperature Thresholds by Mangrove Species




- Leaf damage thresholds were **-5.7 °C** for black mangrove, **-4.1 °C** for red mangrove, and **-3.6 °C** for white mangrove.
- Mortality thresholds were **-6.2 °C** for black mangrove, **-4.4 °C** for red mangrove, and may occur below **-3.8 °C** for white mangrove.

Recovery of Black Mangroves

- Resprout branches of surviving black mangroves, from both stem and base, were positively correlated with leaf damage.
- Resprout was activated at a temperature threshold of **-5.3 °C**.




Freeze Responses by Temperature, Species, and Stratum



- Mortality rate and leaf damage of red mangroves were higher in tall stratum than short stratum, while no difference were found between black mangroves strata.
- Mortality rate and leaf damage of tall red mangrove significantly decreased with temperature.

CONCLUSION

- Temperature thresholds for freeze resistance were quantified for three mangrove species; black mangroves were the most resistant, followed by red and white mangroves.
- All three mangrove species resprouted after freeze damage, with black mangroves being most freeze-resilient.
- Population recovery of red mangroves may primarily depend on growing of well-preserved short trees, despite high freeze mortality and little resprout in tall red mangroves.



References
¹Cavanaugh, K.C., Kellner, J.R., Forde, A.J., Gunter, D.S., Parker, J.D., Rodriguez, W., Feller, I.C., 2014. Poleward expansion of mangroves is a threshold response to decreased frequency of extreme cold events. Proc. Natl. Acad. Sci. U. S. A. 111, 723-727.
²Kaustad, S., Osland, M.J., Devlin, D.J., Proffitt, C.E., Feher, L.C., Armitage, A.R., Day, R.H., Swanson, K.M., Anderson, G.H., Berger, B., 2023. Temperature Thresholds for Leaf Damage from Two Extreme Freeze Events (2018 and 2021) Near the Northern Range Limit of Black Mangroves (Avicennia germinans) in Southeastern North America. Estuaries Coasts, 1-9.
³Ross, M.S., Ruiz, P.L., Sali, J.P., Hanan, E.J., 2009. Chilling damage in a changing climate in coastal landscapes of the subtropical zone: a case study from south Florida. Global Change Biology 15, 1817-1832.

Acknowledgement
 We thank H. Sackles, Y. Serra, and A. Jackson for their help in field data collection. This research was funded by the Nature Coast Biological Station of University of Florida and the United States Geological Survey.