



# Habitat use and movement patterns of the American alligator (*Alligator mississippiensis*) in the Florida Coastal Everglades

Adam E. Rosenblatt<sup>1\*</sup>, Michael R. Heithaus<sup>1</sup>, Frank J. Mazzotti<sup>2</sup>

<sup>1</sup>Florida International University (\*Correspondence: arose007@fiu.edu), <sup>2</sup>University of Florida



## Abstract

American alligators (*Alligator mississippiensis*) are top predators in the Florida Coastal Everglades (FCE), but their movement patterns and role in the ecosystem are largely unknown. Proposed restoration and management efforts in the Everglades are likely to influence both alligator movements and the spatiotemporal patterns of their effects on ecosystem and community dynamics, making studies of the factors influencing alligator movements important at this time. In October 2007, we initiated a study of alligators in the Shark River Slough of Everglades National Park using a tracking technique that has yet to be used on crocodilians: passive acoustic telemetry. Acoustic transmitters attached to 12 alligators have been detected 34,000 times on an array of 42 monitoring stations within the river system from Oct. '07 – Feb. '08. Acoustic transmitters provide temporally detailed information on movements within a monitoring array and over a long duration (potentially years). Our preliminary results show that alligators may remain in localized areas for weeks at a time, but make long-distance moves from the mouth of the river system to the marsh-mangrove ecotone and likely into the marsh. Additionally, some individuals followed a definite pattern in their movements, “bouncing” between fresher upstream areas and the saltier Gulf of Mexico repeatedly, possibly in search of prey. In general, most alligators moved upstream as freshwater inputs decreased, but some individuals remained near the river mouth even when salinities were high. Ultimately, by characterizing alligator movements in relation to spatiotemporal variation in environmental factors we will be able to predict how alligator distributions and populations will respond to planned increases in freshwater flow. Furthermore, movement data combined with dietary and stable isotopic studies may shed light on the possible role of alligators in the redistribution of nutrients throughout the Shark River Slough.

## Background and Objectives

- The ecological role and movements of alligators in coastal estuaries are poorly known
- Due to their sensitivity to high salinities, proposed increases in freshwater flow are predicted to impact alligator movements
- To predict the effects of Everglades restoration on alligators, we are assessing their movement patterns using a novel technology



Fig. 1 – Left to right: an alligator being measured, an acoustic transmitter attached to an alligator, and a monitor ready to be deployed.

## Methods

- Alligators captured using pole and snare technique, then measured (Fig. 1), tagged, and outfitted with an acoustic (n=12) transmitter (Vemco, V16) attached to the tail scutes (Fig. 1)
- Acoustic data logged by underwater array (Fig. 2) of 42 monitoring stations (Vemco, VR2/VR2W)(Fig. 1) and then wirelessly downloaded to a personal computer
- Environmental data gathered continuously by FCE Long Term Ecological Research project and USGS




Fig. 2 – Left: a map of our array of underwater listening devices (positions marked in yellow). Right: a screenshot of the VUE program that stores and plots our acoustic movement data.

## Results

- Acoustic units provided ~20 positions per individual per day over 4 months
- Alligators showed prolonged periods of residency but generally moved upstream as salinities increased (Fig. 3, Fig. 4)
- Some alligators made regular movements to higher salinity waters




Fig. 3 – Left: a typical movement pattern of an acoustically tracked alligator from Nov. '07-Feb. '08. Right: Graph of the typical salinity pattern in the Shark River during the dry season.

**Thanks to:** entire Heithaus lab, all the volunteers, Mike Cherkiss, Brian Jeffery, Dr. Craig Layman, Dr. Evelyn Gaiser, Linda Powell, the FCE LTER group, UF, and FIU. Funding has been provided by FIU, UF, and the FCE LTER project.

## Discussion

- Acoustic tags provided many tightly packed positions (Fig. 2)
- As expected the majority (80%) of tracked alligators have moved away from high salinity areas as the dry season has progressed (Fig. 4), suggesting that increased freshwater inputs planned for Everglades restoration may move alligator habitat seaward, which may have community and ecosystem level impacts




Fig. 4 – Left: a graph of the proportion of time each alligator (from a representative subsample) spent upstream as the dry season progressed. Right: a map of the three river zones used in analysis.

- Regular movements up and downstream by several individuals, including to the river mouth and high salinity waters, suggests that alligators could play a role in upstream nutrient transport in the coastal Everglades. However, future studies of alligator diets and stable isotopic signatures are required to evaluate this possibility
- Movements appear to be directed into the marsh as salinity rises, which may result in alligators bringing marsh nutrients down into the river at the onset of the wet season