Seagrass Non-Cryptic Fish Assemblage Across the Florida Keys National Marine Sanctuary Seascape














Predator Ecology & Conservation Lab

Based at FIU, our team of ecologists, difference for sharks, rays, and reef fish









Seagrass Ecosystems Research Laboratory



FISHSCAPE PROJECT QUESTION:

How much seagrass is necessary to support

reef fish foraging? P.I: Alastair Harborne



This project is funded by the NOAA's National Centers for Coastal Ocean Science, Competitive Research Program under award NA21NOS4780150 to Florida International University.



This project is funded through the South Florida-Caribbean Cooperative Ecosystems Studies Unit

Multi-scale approach

Seascape mapping

Acoustic telemetry to track fish movements

Laboratory experiments measuring energetic costs

Mesocosm Experiment On Mesopredatory fish Foraging Efficiency

Stable Isotope Analyses and Mesopredators diet

Foraging fishes and Predators/Prey assessment with **Baited Remote Underwater Video Stations-BRUVS**





UNIVERSITY



Non- cryptic Fish Community Assemblage With Baited Remote Underwater Video Systems-BRUVS

Objective:

To characterize fish assemblage patterns across different seagrass habitats at three distinct regions within the FKNMS, examining how seascape variables influence the non-cryptic fish assemblage.



Methods- Baited Remote Underwater Video Stations- BRUVS

- Non-invasive. Camera mounted on a frame with a bait arm extending one meter from the camera
- The bait cage at the end contains standardized bait to attract fish into the camera's field of view


Key advantages of BRUVS:

- Records species missed by visual census/trawling (Mesopredators and Upper predators)
- Captures rare species occurrences
- Documents natural behavior
- Allows abundance quantification





Baited Remote Underwater Video Stations




102 BRUVS were deployed in mornings in sites at different seagrass continuity, distances to shore and reefs.

 Units were placed at least 500 meters apart to ensure independent sampling

• Each frame recorded >170 minutes = > 290 hours.



MaxN (Cappo et al., 2003)



The maximum number of individuals of a species visible in a single video frame.

This conservative measure avoids double-counting individuals and provides reliable data for my analyses.




Environmental and Seascape variables



Depth



Temperature



Turbidity (5 levels scale)



Turbidity

5-point qualitative scale based on water color and visibility in the video frames.



2. Clear (blue to blue-teal water with good visibility) 3. Moderate turbidity (blue-green. Moderate visibility)

4. Turbid (Green water with reduced visibility) 5. Very turbid (intense green water with poor visibility)





Environmental and Seascape variables



Depth



Temperature



Turbidity (5 levels scale)




Gravity Index-Fishing pressure



Gravity Index (Cinner et al., 2018)

Human interactions with a reef are a function of the population of a place divided by the squared time it takes to travel to the reefs





Environmental and Seascape variables



Depth



Temperature



Turbidity (5 levels scale)




Gravity Index-Fishing pressure (Cinner et al., 2018)



Distance to Shore



Distance to reefs



Reef connectivity (Buffers of 100m, 250m, 500m)




Seagrass continuity (Florida's Unified Reef Map)



Continuous/Discontinuos Seagrass







Fish Assemblage Analysis

Diversity Metrics

Alpha diversity

Species
accumulation
curves and richness
estimation

Boosted Regression Trees (BRT)


Quantified relative influence (%) for each predictor in Species abundances and diversity

Multivariate Analysis (dbRDA)

Distance-based Redundancy Analysis with Bray-Curtis to identify which factors drive regional differences



Results




- 95 Non-cryptic fish species across FKNMS seagrass habitats
- Species estimation is 120 species
- BRUVS + Visual Census (by FIU Seagrass Ecosystems Research Lab) = 174 species



Figure 1: Species accumulation curve

Diversity Patterns by location




Upper Keys show significantly lower species richness (~9 species) compared to Lower Keys and Middle Keys (~12-13 species, p < 0.001)

Middle Keys exhibit the highest diversity (~2.1 H') with significant differences among regions (p < 0.001)



Top dominant species across regions




- 1. Yellowtail abundance
- Snapper- Highest



- 3. Yellow Jack
- 4. Pinfish -**Dominant in Lower** Keys (~30%).










Frequency of Occurrence of Fish Species in FKNMS Seagrass

Top 10 Most Frequent Species (All Regions Combined)



The Yellowtail Snapper, our most frequently observed species, shows high occurrence rates across all regions


Sphyraena barracuda was the most common predator (60% Frequency of Occurrence)





Other predators and rare species

- 7 Shark species: Bull shark, Nurse shark, Blacktip shark, Caribbean Reef Shark, Great Hammerhead Shark, Bonnethead Shark, Tiger Shark.
- Other Species: Cormorants, Turtles, Dolphins









Boosted Regressions Trees results:

Variable Importance from 100-Iteration Averaged BRT Models

Mean relative influence (%) across 100 independent model runs with relationship direction + positive | - negative | 0 weak/non-linear | - negligible (<3%)

	Meso-Invertivores	20 -	12 +	14 -	19 +	20 -	2	10 -	1	0
Response Variable	Top Predators	7 -	44 +	16 -	8 °	4 °	4	10 -	1	4 -
	Great Barracuda	22 -	21 +	16 -	<u>8</u>	15 +	1	13 +	2	<u>2</u>
	White Grunt	22 -	10 +	17 -	25 +	10 -	4	9 -	2	2_
	Yellowtail Snapper	66 -	12 +	3_	2_	0	14	2_	1	0
	Total Abundance	14 +	29 +	21 -	16 +	12 -	2	4 -	1	1_
	Simpson Diversity	18 +	<u>4</u> -	18 +	12 -	<u>4</u> -	27	13 +	2	1
	Species Richness	16 -	46 +	10 +	9	3 -	12	3 +	1	0

Distance to shore land

Depth ft.

Gravity index

Location Temperature

grass continuity

Relative Influence


> 30 20 10



Boosted Regressions Trees results:

Variable Importance from 100-Iteration Averaged BRT Models

Mean relative influence (%) across 100 independent model runs with relationship direction + positive | - negative | o weak/non-linear | - negligible (<3%)












Distance to shore relationships with the most important species







Yellowtail Snapper

White Grunt

Great Barracuda




Boosted Regressions Trees results:

Variable Importance from 100-Iteration Averaged BRT Models

Mean relative influence (%) across 100 independent model runs with relationship direction





Distance to shore [km]

Dephilit

Distance to Reet In

Location

emperature er

Turbidityleve

Relative

Influence

50

40

30 20


10



Boosted Regressions Trees results:


Variable Importance from 100-Iteration Averaged BRT Models


Mean relative influence (%) across 100 independent model runs with relationship direction + positive | - negative | o weak/non-linear | - negligible (<3%)






Distance to reef the most important for White Grunt







Distance-based Redundancy Analysis (dbRDA)





Take home message:

1. Seagrass habitats across FKNMS support 95 non-cryptic fish species (~174 including cryptic with visual census) High diversity!





2. Distance to shore influence mesopredators abundance





Family **Haemulidae**.

It is a **nocturnal forager** on benthic invertebrates and fish occupying **reef habitats** during the day and **seagrass** at night.

This species holds **economic importance**.



Yellowtail Snapper (Ocyurus chrysurus)

Family Lutjanidae

It exhibits a more **mobile feeding strategy**

Considered an economically valuable species.



Acknowledgments

FISHSCAPE project collaborators

Kirk Gastrich and Global FinPrint for use of the BRUVS

TFEL Lab members





















