

# Citrus: Should We Still Plant Them?

## *The Question of the Hour.*

J.D. Burrow, M.M. Dewdney, and M. Rogers

University of Florida, IFAS, Citrus Research and Education Center

# Outline

- Florida Citrus History
- Citrus Canker
- Citrus Greening
- Homeowner Options for Growing Citrus
- Anything new on the horizon?
- Summary: The answer to the question.



# Florida Citrus History

- 1565-Citrus arrived with Spaniards in St. Augustine
- 1763-First commercial citrus grove established
- 1860-Leesburg was declared the first citrus center of the state
- 1915-First citrus processing plant in America opens
- 1945-Citrus concentrate is developed
- 1989-“Christmas Freeze” practically wiped out the citrus industry north of I-4

# Florida Citrus History

- Florida is closely linked to citrus
- Today, it is a \$9 billion industry

**Timeline of Citrus in Florida**

From the earliest explorers to modern-day pioneers, citrus has been a key ingredient in the rich history of Florida.

Special Thanks to Florida State Archives, USDA, and Florida Southern College's Citrus Archives

Florida Grewer

**Discovery**

- 1513** — Spanish explorer Ponce de Leon, Cuba, introduced citrus to Florida, where it is raised by native Indians.
- 1733** — The first citrus grove in Florida was established by John B. Moore in 1733.
- 1774** — The first references to the shipment of fruit out of Florida appear. The first commercial citrus grove was established in 1774.

**The Early Years**

- 1774** — The first citrus grove in Florida was established by John B. Moore in 1774.
- 1818** — Commercial groves appear on Florida's west coast along the border from 1818 on. The first seedling and transplant into west orange varieties.
- 1870** — General Henry Shelton Sanford comes to the Gulf Coast and plants citrus groves. He is the first to plant large blocks of groves and plant citrus groves. The first citrus grove in the world is established in 1870.
- 1870** — General Henry Shelton Sanford comes to the Gulf Coast and plants citrus groves. He is the first to plant large blocks of groves and plant citrus groves. The first citrus grove in the world is established in 1870.
- 1870** — General Henry Shelton Sanford comes to the Gulf Coast and plants citrus groves. He is the first to plant large blocks of groves and plant citrus groves. The first citrus grove in the world is established in 1870.

**The Growth of Days**

- 1900** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 1900** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 1900** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.

**New Frontiers**

- 1945** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 1945** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 1945** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.

**2005**

- 2005** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 2005** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.
- 2005** — The Florida Citrus Commission is established in Tampa. For more than 80 years, the Florida Citrus Exchange would operate under the name Sanford, Moore and Son.

# Florida Citrus Disease-Pest History

- 1933-Citrus Canker (bacterial disease)
- 1993-Citrus Leafminer (pest)
- 1998-Asian Citrus Psyllid (pest)
- 1986 and 1995 – Citrus Canker returns
- 2005-Citrus Greening (bacterial disease)
- 2010-Citrus Black Spot (fungal disease)
- Future...the possibility of diseases we don't have yet!

# Citrus Greening

# Fact or Fiction

- Huanglongbing
  - HLB
  - Citrus greening
- Spread by an insect



# History

- 1919: First reported in China
- 1921: Reported in the Philippines, but thought it was zinc related
- 1937: In South Africa, thought to be mineral toxicity
- 1941-1955: Most extensive work on greening conducted in southern China






# History

- 1960: Appeared in Thailand
- 1965: Researchers demonstrated HLB was transmissible by grafting and the citrus psyllid
- 1966: Filipino and Indian researchers recognized the similarities between various named diseases




# History

- 1998: Asian citrus psyllid arrived in Florida
- 2004: Disease confirmed in Brazil
- 2005: Disease confirmed in Florida
- 2005 to the present: Disease continues to spread throughout Florida



# United States Locations



Map retrieved from <http://www.martinsaphug.com/learn/maps-2/united-states-and-canada/>

# Importance


- Affects fresh market fruit
- Affects processed fruit
- No cure for the disease






# Biology

- Caused by a bacteria
- Found within the phloem of the tree



**Healthy**



**Infected**

# Biology

- Affects all citrus varieties
- Affects plants in the *Rutaceae* family
- Affects box orange and orange jasmine



*Murraya paniculata*  
(orange jasmine)

# Biology

- Some plants host the vector only
- Some plants host the vector and are susceptible for greening

Known Host Plants of Huanglongbing (HLB) and Asian Citrus Psyllid

Plant Name	<i>Liberibacter asiaticus</i> Huanglongbing	<i>Diaphorina citri</i> Citrus Psyllid
<i>Aegle marmelos</i> (L.) Corr. Serr.: baol, Bengal quince, golden apple, bela, mulva		X
<i>Aeglopsis chevalieri</i> Swingle: Chevalier's aeglopsis	X	X
<i>Aegle gabonensis</i> (Swingle) Engl.: Gabon powder-flask		X
<i>Aegle paniculata</i> (Schum.) Engl.: Nigerian powder-flask		X
<i>Azolania missouriensis</i> (Wall. ex Wright) Oliv.: see <i>Famburus missouriensis</i>	X	X
<i>Azolania monophylla</i> (L.) Corr.: Indian atlantia		X
<i>Balsanocitrus dawsoni</i> Stapf: Uganda powder-flask	X	X
<i>Burkillanthus malaccensis</i> (Ridl.) Swingle: Malay ghost-lime	X	
<i>Calodendrum capense</i> Thunb.: Cape chestnut	X	
= <i>Citroncitrus weddeleri</i> J. Ingram & H. E. Moore: citrange	X	
<i>Citropsis gileniana</i> Swingle & M. Kellerman: Gillet's cherry-orange		X
<i>Citropsis schweinfurthii</i> (Engl.) Swingle & Kellerman: African cherry-orange		X
<i>Citrus amblycarpe</i> (Hassk.) Ochs.: djerook leemo, djeruk-limau	X	
<i>Citrus aurantiifolia</i> (Christm.) Swingle: lime, Key lime, Persian lime, lima, limón agrio, limón ceuti, lima majicana, limero	X	X
<i>Citrus aurantium</i> L.: sour orange, Seville orange, bigarade, marmalade orange, naranja agria, naranja amarga		X
<i>Citrus depressa</i> Hayata: shikishiwashita, shikishwasita, sequease	X	
<i>Citrus grandis</i> (L.) Osbeck: see <i>Citrus maxima</i>	X	
<i>Citrus hassaku</i> hort. ex Tanaka: hassaku orange	X	
<i>Citrus hystrix</i> DC.: Mauritius papada, Kaffir lime	X	X
<i>Citrus ichangensis</i> Swingle: Ichang papada	X	
<i>Citrus jambhiri</i> Lushington: rough lemon, jambhiri-orange, limón rugoso, rugoso	X	X
<i>Citrus junos</i> Sieb. ex Tanaka: siang cheug, yumi	X	
<i>Citrus kabachi</i> hort. ex Tanaka: this is not a published name; could they mean <i>Citrus kinokuni</i> hort. ex Tanaka, kishu nikan?	X	
<i>Citrus limon</i> (L.) Burm. f.: lemon, limón, limonero	X	X
<i>Citrus × limonia</i> Osbeck: potted orange, Rangpur lime, mandarin lime, lemandarin, limón de Canton, limón grande	X	
<i>Citrus maxima</i> (Burm.) Merr.: pummelo, pomelo, shaddock, pomelo-sinuso, toronja	X	X
<i>Citrus medica</i> L.: citron, cidra, cidro, toronja		X
<i>Citrus meyeri</i> Tan.: Meyer lemon, dwarf lemon		X
<i>Citrus × nobilis</i> Lou.: king mandarin, tangor, Florida orange, King-of-Siam	X	X
<i>Citrus oto</i> hort. ex Tanaka: mandarin?	X	
<i>Citrus × paradisi</i> Macfad.: grapefruit, pomelo, toronja	X	X
<i>Citrus reticulata</i> Blanco: mandarin, tangerine, mandarina	X	X
<i>Citrus sinensis</i> (L.) Osbeck: sweet orange, orange, naranja dulce, naranja	X	X
<i>Citrus sunki</i> (Hayata) hort. ex Tanaka: sour mandarin, suki mandarin	X	
<i>Citrus unshiu</i> (Mack.) Marc.: satsuma orange, satsuma mandarin		X
<i>Clausena anisum-olens</i> (Blanco) Merr.: anis (Philippines)		X
<i>Clausena excavata</i> Burm. f.: clausena		X
<i>Clausena indica</i> Oliv.: clausena	X	X

Plant Name	<i>Liberibacter asiaticus</i> Huanglongbing	<i>Diaphorina citri</i> Citrus Psyllid
<i>Clausena lansium</i> (Lour.) Skeels: wampee, wampee	X	X
<i>Cuscuta australis</i> R. Br.: dodder, cuscuta	X	
<i>Eremocitrus glauca</i> (Lindley) Swingle: Australian desert lime		X
<i>Fortunella</i> spp.: kumquat	X	
<i>Fortunella crassifolia</i> Swingle: Meirwa kumquat		X
<i>Fortunella margarita</i> (Lour.) Swingle: Nagami kumquat, oval kumquat		X
<i>Fortunella polyandra</i> (Ridley) Tanaka: Malayan kumquat		X
<i>Limonia acidissima</i> L.: Indian wood apple	X	X
<i>Merrillia caloxylon</i> (Ridley) Swingle: flowering merrillia		X
<i>Microcitrus australasica</i> (F. J. Muell.) Swingle: finger-lime	X	X
<i>Microcitrus australis</i> (Planch.) Swingle: Australian round-lime		
<i>Microcitrus papuana</i> H. F. Winters: desert-lime		
<i>Microcitrus 'Sydney'</i> : see <i>Microcitronella</i>		X
= <i>Microcitronella 'Sydney'</i> : fanstrimelin		X
<i>Murraya exoniata</i> L.: see <i>Murraya paniculata</i>		X
<i>Murraya koenigii</i> (L.) Sprengel: curry leaf	X	X
<i>Murraya paniculata</i> (L.) Jack: Lakeview, orange-jasmine, Chinese-box, naranyo jarama	X	X
<i>Naranyo crenulata</i> (Rorb.) Nicholson: naranyo		X
<i>Pamburus missouriensis</i> (Wall. ex Wright) Swingle: pamburus		X
<i>Poncirus trifoliata</i> (L.) Raf.: trifoliata orange	X	X
<i>Severinia buxifolia</i> (Poir.) Ten.: Chinese box orange	X	X
<i>Swinglea glutinosa</i> (Blanco) Merr.: tabog	X	X
<i>Toddalia asiatica</i> (L.) Lam.: orange climber		X
<i>Toddalia lanceolata</i> Lam.: toddalia	X	
<i>Triphasia trifolia</i> (Burm. f.) P. Wilson: trifoliata limeberry	X	X
<i>Tigridis lanceolata</i> (Lam.) G. Don: white tougwood		X
<i>Zanthoxylum fagara</i> (L.) Sarg.: wild lime, lime prickly-ash		X

Updated 12/22/2008

# Bacterium


- *Candidatus Liberibacter asiaticus*
- Gram negative
  - Defines type of bacteria
- Phloem limited
- Fastidious bacterium
  - Cannot grow in culture
- Reproduces/multiplies in both the psyllid and the tree




Photo Credit: Huanglongbing: A Destructive, Newly-Emerging, Century-Old Disease of Citrus, J.M. Bové



LAS must be present and associated with all diseased plants examined.




A sample is taken from the infected plant material. LAS must be isolated from the plant material and grown in pure culture.



*This step has yet to be achieved.*


A sample of the possible disease causing organism from a pure culture is inoculated into healthy plant material of the same species or variety from which it was originally taken.




# Koch's Postulates

The procedure required to show that *Ca. Liberibacter asiaticus* (LAS) causes Huanglongbing

The bacterium must be reisolated in pure culture from the inoculated host and the new culture must have the same characteristics as seen in the original pure culture.



A sample needs to be taken from the inoculated host.



The healthy tree must produce the same symptoms displayed by the original tree.

# Spread

- Asian citrus psyllid
- Grafting with infected bud wood
- Is not seed transmissible
- **NOT** by contact, tools or equipment

# Asian Citrus Psyllid

- 5 nymphal stages
- Ten generations per year
- Life cycle between 15-47 days
- Egg to adult in two weeks at 75-80°F



# Asian Citrus Psyllid

- Psyllids fly or are carried by the wind to new plants
- Psyllids feed on an infected tree and then transmit the bacteria to healthy trees



# Asian Citrus Psyllid Damage

- Spreads the greening bacteria
- Nymphs produce a waxy secretion
- Notching on leaves





# Leaf Symptoms

- Blotchy mottle patterns
  - Mature leaves
  - Asymmetrical pattern
  - Inside or outer edges of canopy
  - Pattern will appear on both sides of the leaf




# Leaf Symptoms

- Blotchy mottle patterns
- Yellow veins
  - Not a definite symptoms of HLB, but one should inspect the tree more closely if found
  - Found on young and mature leaves

# Leaf Symptoms

- Blotchy mottle patterns
- Yellow veins
- Vein corking
  - Raised veins with a corky appearance
  - Found on mature leaves





# Commonly Mistaken for HLB

- Broken limb
- Foliar symptoms of trees with foot rot (Phytophthora)



# Commonly Mistaken for HLB

- Insect damage
- Herbicide/Chemical damage





Don't forget to  
look at both  
sides!



# Nutrient Deficiencies vs. HLB

- Zinc
  - Small and narrow leaves with yellow mottle on green background
- Iron
  - Green veins on a light yellow to white colored leaf



Mongi Zekri

# Nutrient Deficiencies vs. HLB

- Manganese
  - Dark green veins with a lighter green background
- Magnesium
  - Inverted 'V' pattern



Mongi Zekri





Tom Obreza



# Identifying a Leaf Sample

- Circle areas on opposite sides of the midvein. Are they the same on both sides?
  - Nutrient deficiencies are symmetrical and HLB symptoms are asymmetrical
- Look at the other side of the leaf



# Fruit Symptoms

- Unmarketable, bitter fruit
- Cannot be used for fresh
- Can be used in process, but may have a flavor consequence



# Internal Fruit Symptoms

- Yellow stain beneath the calyx button
- Curved central core
- Aborted seeds





# External Fruit Symptoms

- Lopsided
- Misshapen
- Small
- Does not color properly





# Tree Symptoms

- Leaf and fruit drop
- Yellow shoot
- Severely infected trees
  - Stunted
  - Sparse foliage
  - Twig dieback
- Off-season bloom



# Unsure-is it greening or not?

- Three options for testing
  - Southwest Florida Research and Education Center, Immokalee
  - Division of Plant Industry, Gainesville
  - Plant Disease Clinic, Gainesville

# What to do if a tree has greening

- No known cure once a tree is infected
- You cannot prune away an HLB infection
- Tree removal is the only known effective control
- Trees usually decline within 3-5 years of infection and ultimately die

# Homeowner Options

- Prevention
- Tree Removal and replant
- Refuse to remove tree
- Remove ornamental hosts
- Alternative crops

# Option #1: Prevention

- Maintain a healthy tree
- The source of spread is the psyllid; therefore, prevention of the psyllid is key!
- You cannot prevent psyllids flying onto your citrus trees
- Limited chemicals for homeowner usage

# Option #1: Chemical Prevention

- Non-systemic (foliar applied)
- Systemic (soil applied and taken up by roots)

**Remember - the label is the law!**

# Option #1: Chemical Prevention

- Non-systemic
  - Horticultural oil
  - Malathion
  - Carbaryl
- Systemic
  - Imidacloprid

# Option #1: Organic Prevention

- Horticultural mineral oil
- Neem oil

**Remember - the label is the law!**



# Option #1: Organic Prevention

- Kaolin
  - Clay product
  - White covering on the tree
    - Photosynthesis???
    - Reduces heat stress; therefore, maintaining photosynthesis
    - Improves productivity
    - Reduces sunburn (mandarin)
  - Will need to purchase from specialty garden stores



S. L. Lapointe, U.S. Horticultural Research Lab, Ft. Pierce, FL

# Option #2: Tree Removal and Replant

- A psyllid won't feed on a dead tree
- After removal of the tree
  - Burn
  - Put in lawn waste



# Option #2: Tree Removal and Replant

- When removing an infected tree be sure to kill the stump with a herbicide to prevent sprouting
  - Sprouts will contain the bacteria and be a source of inoculum





# Option #2: Tree Removal and Replant

- If other trees nearby, some roots will graft together-be careful applying a herbicide!
- You can replant in the same area, but should wait depending on the chemical used



# Option #2: Tree Removal and Replant

- No regulated time period to replant
- Be cautious and alert when buying a citrus tree
- Citrus trees are more scarce and more expensive than they used to be – buying one is an INVESTMENT!
- Buy from a reputable local garden center or nursery who is knowledgeable about citrus



# Option #2: Tree Removal and Replant

- DO NOT buy a tree without a tag!
  - FDACS Rule 5B-62.020(3)
    - All retail citrus trees must be tagged with information to identify the variety of the rootstock and scion stock and producing nursery. Each individual tree shall be identified with a slip-on label displaying the following information:
      - a. The producing nursery's certificate of nursery registration number... It is not necessary to include the name of the producing nursery on the label.
      - b. The variety name, rootstock and month and year acquired.



# Option #3: Refuse to remove tree

- Tree remains a source for psyllids to feed and continue to spread the bacteria
- It will become unproductive!



# Option #3: Refuse to remove tree

- Nutrient programs?
  - Will not cure the tree, only prolong the life

**Growing citrus in the dooryard is no longer a simple task.**

**Options 4 and 5 will assist in a different way.**

# Option #4: Remove ornamental hosts

- Remove host plants



*Murraya paniculata*  
(orange jasmine)



# Option #5: Alternative Crops

- Peaches
- Blueberries
- Pomegranates
- Persimmons
- Chestnuts



# Alternative Crops-Considerations

- Pick something you like
- Ornamental flowers, but not edible
- Maintenance
- Varmits (squirrels, raccoons, deer, etc.)



# Anything new on the horizon?

- Research is on-going
- Research is wide-ranging
- Research takes time!



# Summary

- The answer to the question
- Ask yourself these questions
  - Do I have time to take care of my tree(s)?
  - Do I have the financial means to purchase the necessary products to prevent citrus greening?
  - If I answered yes to the above questions, will I make the commitment?

**Growing citrus today is an investment of time, money, and resources.**

**Are you willing or are you unable?**



***There are many opinions on whether residents should or should not grow citrus, but in reality, the decision is left to the stewardship of the homeowner and their willingness to be an asset to the citrus industry.***

\*According to the Merriam-Webster Dictionary, stewardship is defined as the conducting, supervising, or managing of something; the careful and responsible management of something entrusted to one's care

**Any questions?**