

# Diet and Selectivity of the Purple Swamphen in South Florida

Corey T. Callaghan & Dale E. Gawlik

**FAU** ENVIRONMENTAL  
SCIENCE PROGRAM

Charles E. Schmidt College of Science  
Florida Atlantic University

# Background

## -Nonnative avian species in Florida

- ▶ Florida is heavily populated with nonnative avian species
- ▶ In 1992, there were 146 exotic avian species and by 2014, there are about 225




# Background

## -Potential Problems

- ▶ The degree to which the Purple Swamphen poses a threat to native fauna in Florida is currently unknown
- ▶ Competition
  - ▶ Aggression between Great Blue Herons and swamphens
  - ▶ Have preyed upon Black Swan eggs in Australia
  - ▶ Have been observed carrying a presumed Black-necked Stilt chick in Florida
- ▶ Altering vegetation structure and composition
  - ▶ Multiple nesting platforms
  - ▶ Large body size
- ▶ The threat largely depends on the degree of diet and habitat overlap with native species

# Background

## -Study Species


- ▶ Purple Swamphen 

(*Porphyrio porphyrio*)

- ▶ *Rallidae* family

- ▶ secretive birds

- ▶ Closely related to

- ▶ Purple Gallinule 

- ▶ Common Gallinule


- ▶ American Coot




# Background

## -Study Species

### ▶ Large global range



# Background




## -Introduction to Florida

- ▶ First discovered in South Florida in 1996
  - ▶ escaped from an aviculturist
- ▶ By 1999, 135 swamphens were counted in the vicinity of the initial discovery
- ▶ In Florida, chicks have been observed during all months except June, September, November, and December
- ▶ Swamphens can move more than 300 km to colonize new habitats
- ▶ In October 2006, an eradication program was attempted
  - ▶ over 3,100 swamphens were culled

# Background

-Population expansion



(Pranty 2012)


# Background

## -Current diet information

- ▶ In their native home Australia, swamphens are known to eat predominantly herbaceous materials from the families *Graminae* (59%), *Cyperaceae* (17%), and *Hydrocharitaceae* (11%)
- ▶ Predominantly herbivores, but individuals may feed on small animal prey such as: mollusks, leeches, crabs, fish, frogs, birds and their eggs, and rodents
- ▶ Little is known about the diet of Purple Swamphens in Florida
  - ▶ swamphens' stomachs in the STAs were bursting with rice grains

# Objectives

- ▶ (1): Quantify the diet of the Purple Swamphen (*Porphyrio porphyrio*)
- ▶ (2): Identify any selectivity the swamphens may demonstrate
- ▶ (3): Compare morphological measurements of the swamphens among three study sites



# Hypotheses

- ▶ (1): Swampheens will predominantly be herbivorous as they are throughout most of their range

- ▶ (2): Swampheens select for *Eleocharis spp.*




Image credit: Tyler Beck

# Methodology

## -Collection


- ▶ Three study sites
- ▶ Sample size
  - ▶ WCA2B: 32
  - ▶ STA1W: 27
  - ▶ Lake Okeechobee: 24



# Methodology

## -Objective (1): Diet

- ▶ Contents removed from proventriculus, gizzard, and crop
  - ▶ Stored in 70% ethanol
- ▶ A macroscopic and microscopic level of sorting and identification
- ▶ Reference collection utilized
- ▶ Stepwise process
  - ▶ Macroscopic sorting
  - ▶ Random sampling and identification of the remaining contents (homogenate)
  - ▶ Dry weight of the items



# Methodology


## -Objective (1): Diet

- ▶ Statistical Analysis: MDS and ANOSIM using PRIMER
  - ▶ Multi-dimensional Scaling
    - ▶ Visualization of similarity/dissimilarity
  - ▶ Analysis of Similarity
    - ▶ Compares variation and composition among sites

# Methodology

## -Objective (2): Selectivity

- ▶ Vegetation sampling area was defined
- ▶ Nested plot design at random points in which each point represents the Northeast corner a priori
  - ▶ 5m x 5m, 3m x 3m, and 1m x 1m
- ▶ Utilized a modified Braun-Blanquet scale to estimate the percent cover at each point



Braun-Blanquet scale	Range of cover (%)
5	75-100
4	50-75
3	25-50
2	5-25
1	<5



# Methodology

## -Objective (2): Selectivity

- ▶ Chesson's Index:  $\alpha_i = \frac{r_i/p_i}{\sum r_i/p_i}$ ,  $i = 1, \dots, m$ 
  - ▶  $\alpha_i$  is the selectivity index for prey type  $i$
  - ▶  $r_i$  is the relative abundance of prey type  $i$  in the swamphen's stomach
  - ▶  $p_i$  is the percent of prey type  $i$  in the environment calculated from the vegetation surveys
  - ▶  $m$  is the number of prey types available in the environment
- ▶ Chesson's index is interpreted by relating  $\alpha_i$  to  $1/m$ 
  - ▶ Random feeding:  $1/m = \alpha_i$
  - ▶ Selection:  $\alpha_i > 1/m$
  - ▶ Avoidance:  $\alpha_i < 1/m$
- ▶ Assumptions
  - ▶ Prey abundance is large
  - ▶ Ability to consume a species is equal
  - ▶ Plants detected by us are also detected by swamphens

# Methodology


## -Objective (3): Morphology




- ▶ Morphological measurements taken before dissection:
  - ▶ Body mass
  - ▶ Tarsus length
  - ▶ Bill length to gape
  - ▶ Wing chord
  - ▶ Exposed culmen
  - ▶ Tail length
  - ▶ Bill width
  - ▶ Pectoral score
  - ▶ Bill depth
- ▶ Statistical Analysis: MDS and ANOSIM using PRIMER




# Findings

- ▶ **(1): Quantify the diet of the Purple Swamphen (*Porphyrio porphyrio*)**
- ▶ (2): Identify any selectivity the swamphens may demonstrate
- ▶ (3): Compare morphological measurements of the swamphens among three study sites





$R=0.525, p<0.1$





2D Stress: 0.135

*Location*

▲ STA1W

▼ LKO

■ WCA2B




# Findings

- ▶ (1): Quantify the diet of the Purple Swamphen (*Porphyrio porphyrio*)

- ▶ **(2): Identify any selectivity the swamphens may demonstrate**


- ▶ (3): Compare morphological measurements of the swamphens among three study sites



◆ Gulf Coast spikerush (*Eleocharis cellulosa*)


▲ *Panicum spp.*

■ Sawgrass (*Cladium jamaicense*)



# Findings


- ▶ (1): Quantify the diet of the Purple Swamphen (*Porphyrio porphyrio*)
- ▶ (2): Identify any selectivity the swamphens may demonstrate
- ▶ **(3): Compare morphological measurements of the swamphens among three study sites**





$R = 0.164, p < 0.1$

2D Stress: 0.174



# Conclusions

- ▶ Objective (1): Swampheens are mainly herbivores and eat predominantly Gulf Coast spikerush
- ▶ Objective (2): They show a strong preference for Gulf Coast spikerush in WCA2B
- ▶ Objective (3): The largest birds were collected from STA1W



# Implications

- ▶ Swamphen diets were predominantly herbivorous and were more specialized than the literature suggested because of a strong preference for spikerush
  - ▶ Adaptive flexibility hypothesis
- ▶ Potential impacts to native species likely depends on the degree to which they are dependent on spikerush and the degree of selection of spikerush by swamphens
- ▶ High selectivity is not likely to limit the expansion of swamphens, because of the plant's prevalence throughout Florida and the southeastern U.S.

# Acknowledgements

- ▶ Florida Fish and Wildlife Commission provided the funding and collaboration of this work
- ▶ Lab-mates and technicians in the Gawlik Lab



Questions?

