


# Assessment of Chesapeake Climate Change Using A Suite of Airshed, Watershed, and Estuary Models

NCER Albuquerque, NM

April 18, 2024

Low Linker (EPA-CBPO), Gopal Bhatt (Penn State), Richard Tian (UMCES), Jesse Bash (EPA-ORD), Carl F. Cerco (Attain), Isabella Bertani (UMCES), Joseph Zhang (VMS), Xun Cai (VIMS), Gregorio Toscano Pulido (MSU)



*linker.lewis@epa.gov*






# The CBP Climate Change Assessment


### Airshed Model




### Watershed Model





### Tidal Water Quality Model



### Criteria Assessment Procedures




Nutrient and sediment targets for CBP State and Federal Partnership







# Overview of Bay Designated Uses



A. Cross Section of Chesapeake Bay or Tidal Tributary




B. Oblique View of the "Chesapeake Bay" and its Tidal Tributaries






# Elements of Chesapeake Water Quality Climate Risk Assessment





# Components of Climate Change Effect on Tidal Hypoxia



# Approaches, Methods, and Findings from the Watershed



Chesapeake Bay Program  
*Science, Restoration, Partnership*

# An ensemble of GCM projections from BCSD CMIP5<sup>[1]</sup> was used to estimate 1995-2025 temperature change.

?

Data [?] Unavailable [?]

GCM [?] Used [?]

Selection [?] Updated [?]

Updated [?] Ensemble [?] members [?]		
ACCESS1-0 [?]	FGOALS-g2 [?]	IPSL-CM5A-LR [?]
BCC-CSM1-1 [?]	FIO-ESM [?]	IPSL-CM5A-MR [?]
BCC-CSM1-1-M [?]	GFDL-CM3 [?]	IPSL-CM5B-LR [?]
BNU-ESM [?]	GFDL-ESM2G [?]	MIROC-ESM [?]
CanESM2 [?]	GFDL-ESM2M [?]	MIROC-ESM-CHEM [?]
CCSM4 [?]	GISS-E2-H-CC [?]	MIROC5 [?]
CESM1-BGC [?]	GISS-E2-R [?]	MPI-ESM-LR [?]
CESM1-CAM5 [?]	GISS-E2-R-CC [?]	<b>MPI-ESM-MR [?]</b>
CMCC-CM [?]	HadGEM2-AO [?]	MRI-CGCM3 [?]
CNRM-CM5 [?]	HadGEM2-CC [?]	NorESM1-M [?]
CSIRO-MK3-6-0 [?]	HadGEM2-ES [?]	<b>31 member ensemble</b>
EC-EARTH [?] [?]	INMCM4 [?]	

[1] BCSD – Bias Correction Spatial Disaggregation;  
[1] CMIP5 – Coupled Model Intercomparison Project 5



Reclamation, 2013. 'Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections: Release of Downscaled CMIP5 Climate Projections, Comparison with preceding Information, and Summary of User Needs', prepared by the U.S. Department of the Interior, Bureau of Reclamation, Technical Services Center, Denver, Colorado. 47pp.

Source: Kyle Hinson, VIMS



# For the 2025 Climate Change Estimate:

The trends in annual precipitation on a county level were developed through the application of PRISM data and analysis provided and recommended by Jason Lynch, EPA, and Karen Rice, USGS. The annual PRISM dataset for the years 1927 to 2014 (88 years) were used in for the regression trend analysis. For the analysis PRISM data were first spatially aggregated for each Phase 6 land segments. The Phase 6 land segments typically represent a county. For each land segment a simple linear trend was fitted to the annual rainfall dataset.




Annual rainfall volumes for the 88-year period linear regression lines are shown in red for the two land segments (counties) – (a) Centre County in Pennsylvania and (b) District of Columbia. The values for the slope of the regression lines, and the corresponding 30-year projections in the rainfall volume (1995 to 2025) are also shown.





# Assessment of Influence of 2025 Climate Change in the Watershed




## Projections of rainfall increase using trend in 88-years of annual PRISM<sup>[1]</sup> data

### Change in Rainfall Volume 2021-2030 vs. 1991-2000

2025 Rainfall Projection (percent change)

- 0.7 - 1.0
- 1.1 - 1.5
- 1.6 - 2.0
- 2.1 - 2.5
- 2.6 - 3.0
- 3.1 - 3.5
- 3.6 - 4.0
- 4.1 - 4.5
- 4.6 - 5.0
- 5.1 - 5.5
- 5.6 - 6.0



Major Basins	PRISM Trend
Youghiogheny River	2.1%
Patuxent River Basin	3.3%
Western Shore	4.1%
Rappahannock River Basin	3.2%
York River Basin	2.6%
Eastern Shore	2.5%
James River Basin	2.2%
Potomac River Basin	2.8%
Susquehanna River Basin	3.7%
<b>Chesapeake Bay Watershed</b>	<b>3.1%</b>

[1] Parameter-elevation Relationships on Independent Slopes Model




# 1940-2014 streamflow trends based on observations

Chesapeake Bay Program  
Science, Restoration, Partnership

The study analyzed USGS GAGES-II data for a subset of Hydro-Climatic Data Network 2009 (HCDN-2009).


Annual Average Streamflow in the United States, 1940-2014



More than 50% decrease    20% to 50% decrease    20% decrease to 20% increase    20% to 50% increase    More than 50% increase

Data source: USGS (U.S. Geological Survey), 2016. Analysis of data from the National Water Information System. Accessed May 2016.

For more information, visit U.S. EPA's "Climate Change Indicators in the United States" at [www.epa.gov/climate-indicators](http://www.epa.gov/climate-indicators).



Karen C. Rice, Douglas L. Moyer, and Aaron L. Mills, 2017. Riverine discharges to Chesapeake Bay: Analysis of long-term (1927 - 2014) records and implications for future flows in the Chesapeake Bay basin *JEM* 204 (2017) 246-254

USGS station ID	Precipitation		Discharge	
	Slope	p-value	Slope	p-value
04252500	0.0007	<b>0.0011</b>	0.0021	<b>&lt;0.0001</b>
01512500	0.0008	<b>0.0007</b>	0.0016	<b>0.0028</b>
01503000	0.0007	<b>0.0022</b>	0.0013	<b>0.0181</b>
01531000	0.0006	<b>0.0219</b>	0.0018	<b>0.0030</b>
01531500	0.0007	<b>0.0044</b>	0.0016	<b>0.0029</b>
01532000	0.0006	<b>0.0374</b>	0.0015	<b>0.0330</b>
01534000	0.0005	<b>0.0497</b>	0.0015	<b>0.0120</b>
01550000	0.0005	<b>0.0493</b>	0.0019	<b>0.0015</b>
01543000	0.0004	0.1000	0.0018	<b>0.0058</b>
01545500	0.0004	0.0953	0.0017	<b>0.0026</b>
01536500	0.0006	<b>0.0078</b>	0.0016	<b>0.0027</b>
01551500	0.0005	0.0612	0.0017	<b>0.0017</b>
01439500	0.0005	0.0972	0.0007	0.1661
01541500	0.0003	0.2357	0.0017	<b>0.0017</b>
01540500	0.0006	<b>0.0111</b>	0.0016	<b>0.0023</b>
01541000	0.0004	0.0985	0.0016	<b>0.0021</b>
01567000	0.0004	0.1577	0.0011	<b>0.0250</b>
01570500	0.0005	<b>0.0260</b>	0.0013	<b>0.0088</b>

North-South Split

01562000	0.0004	0.1693	0.0007	0.2082
01638500	0.0004	0.1150	0.0008	0.1026
01608500	0.0004	0.1725	0.0010	0.0833
01636500	0.0005	0.1245	0.0008	0.0624
01606500	0.0003	0.1958	0.0009	0.1108
01668000	0.0006	0.0794	0.0004	0.4727
02035000	0.0003	0.2653	-0.0001	0.8243
02019500	0.0002	0.4333	0.0003	0.4836
03488000	0.0003	0.2480	0.0006	0.2841

U.S. Environmental Protection Agency. 2016. Climate change indicators in the United States, 2016. Fourth edition. EPA 430-R-16-004. [www.epa.gov/climate-indicators](http://www.epa.gov/climate-indicators).


Annual average percent change were calculated using Sen slope (Helsel and Hirsch, 2002).

Lins, H.F. 2012. USGS Hydro-Climatic Data Network 2009 (HCDN-2009). U.S. Geological Survey Fact Sheet 2012-3047. <https://pubs.usgs.gov/fs/2012/3047>.

Helsel, D.R., and R.M. Hirsch. 2002. Statistical methods in water resources. Techniques of water resources investigations, Book 4. Chap. A3. U.S. Geological Survey. <https://pubs.usgs.gov/twri/twri4a3>.



# Trends in Observed Rainfall Intensity



**Observed changes in rainfall intensity in the Chesapeake region over the last century. The equal allocation distribution (blue) is contrasted with the distribution obtained based on observed changes (red).**




Chesapeake Bay Program  
Science, Restoration,  
Partnership

# Chesapeake Bay Watershed Annual Change in Temperature

Degrees Celsius

2025 - RCP 4.5




1.1°C Increase  
in Annual  
Temperature

0 35 70 140 210 280 Miles



Degrees Celsius

2050 - RCP 4.5



1.94°C Increase  
in Annual  
Temperature







0 35 70 140 210 280 Miles






# Temperature trends for the six CBP states

Annual temperature for 1895 to 2015 are shown.



Approx. increases over the last 30 years based on the trend line are shown.



NOAA National Climatic Data Center  
<https://www.ncdc.noaa.gov/temp-and-precip/state-temps/>



# Elements of 2025 Climate Change (1995-2025)


**Air-temperature  
increase: 1.06 °C**



↑  
**Sea Level  
Rise:  
0.22m**

**Open boundary:  
Temperature: +0.95 °C;  
Salinity: +0.18 psu  
(Thomas et al., 2017)**

- Flow  
→ 2.4% Increase
- Nitrogen Load  
→ 2.6% Increase
- Phosphorus Load  
→ 4.5% Increase
- Sediment Load  
→ 3.8% Increase




**Phase 6 Watershed Model**

**Model: CH3D-ICM  
400m-1km Resolution**



# Elements of 2035 Climate Change (1995-2035)

**Air-temperature  
increase: 1.39 °C**



Flow  
→  
3.7% Increase

Nitrogen Load  
→  
4.7% Increase

Phosphorus Load  
→  
9.9% Increase

Sediment Load  
→  
8.5% Increase



↑  
**Sea Level  
Rise:  
0.31m**

**Open boundary:  
Temperature: +1.32 °C;  
Salinity: +0.25 psu  
(Thomas et al., 2017)**


**Phase 6 Watershed Model**

**Model: CH3D-ICM  
400m-1km Resolution**




# Estimates of Climate Only and Climate and Land Use


### Marginal Differences in Freshwater Delivery




### Marginal Differences in Sediment Delivery



### Marginal Differences in Nitrogen Delivery



### Marginal Differences in Phosphorus Delivery



2025

2035

2045

2055

2025

2035

2045

2055

Grey bar = climate only    Black bar = Climate and Land Use




# Approaches, Methods, and Findings from the Tidal Bay



Chesapeake Bay Program  
*Science, Restoration, Partnership*

# Summer (Jun.-Sep.) Hypoxia Volume (<1 mg/l) 1991-2000 in the Whole Bay Under 2025 WIP3 Condition







# Bottom DO Change: 1995 to 2025

Keeping all other factors constant, sea level rise and increased watershed flow reduce hypoxia in the Bay, but the predominant influence are the negative impacts of increased water column temperature.


## Sea Level Rise




## Watershed Flow



## Increased Temp.





## All Factors





# Climate Target Loads in Perspective




Overall, the CBP found that a target load of 5 million pounds nitrogen and 0.6 million pounds phosphorus will be sufficient to offset 30 years of climate change in the Chesapeake Bay.

Model load reduction estimates from CAST-2019 (current version of the CBP watershed model)

# ATLAS 14 vs OBSERVED



Comparison of the observed 24-hour rainfall events in the Negley Run Watershed (PA) from 2003-2018 (total) to Atlas 14 estimates (expected)




# CLIMATE CHANGE-INFORMED IDF CURVES

Data Tool: <https://midatlantic-idf.rcc-acis.org/>


Webinar: <https://chesapeakestormwater.net/events/projected-chesapeake-idf-curves/>



# Chesapeake Bay Model Segmentation Viewer



# Chesapeake Bay Model Segmentation Viewer







# Conclusions:

---

Climate change is a multigenerational challenge for the CBP and is a force multiplier for headwinds to the Chesapeake restoration.

However, the CBP is working on management practices that are effective counters to climate change such as:

- The design and accelerated adoption of stormwater management practices appropriately designed for increased rainfall volumes and intensities that are expected in the future for all counties in the Chesapeake watershed.
- Examination of the top tier agriculture and urban BMPs that are most vulnerable to future climate risk, with an emphasis on practices that could be adapted to become more resilient to future climate conditions of increased rainfall intensities and volumes.
- A quantification of the co-benefits of BMPs that mitigate future climate risk.
- Findings in JAWRA Featured Collection *Influence of Climate Change on Chesapeake Bay Water Quality*.

**The climate change risk to the Chesapeake's living resource-based water quality standards can be effectively managed and the CBP is actively addressing the challenge.**